



**Delstrigo**<sup>®</sup>

**doravirine/lamivudine/  
tenofovir disoproxil fumarate**  
100 mg/300 mg/300 mg tablets

*When life brings change,  
what inspires me?*

**MY  
CHOSEN  
FAMILY**

Model living with HIV

**WHAT IS DELSTRIGO?**

DELSTRIGO is a complete, one-pill prescription HIV medicine used to treat HIV-1 infection in adults who have not received HIV-1 medicines in the past, or to replace their current HIV-1 medicines for people whose healthcare provider determines that they meet certain requirements. HIV-1 is the virus that causes Acquired Immune Deficiency Syndrome (AIDS).

**IMPORTANT SAFETY INFORMATION**

**Worsening of hepatitis B virus infection (HBV).** If you have both Human Immunodeficiency Virus-1 (HIV-1) and HBV and stop taking DELSTRIGO, your HBV may suddenly get worse. Do not stop taking DELSTRIGO without first talking to your healthcare provider, as they will need to monitor your health. Your healthcare provider should test you for HBV infection before you start treatment with DELSTRIGO.

# READY FOR CHANGE

**W**hen our lives change, we can change too. By finding fresh inspiration in the people around us. New joy in the things that complete us. And new choices in the treatments available to us. If you are considering your HIV-1 treatment options, it may be time to learn about DELSTRIGO.

**LIFE CHANGES.  
STAY INSPIRED.**

## HHS TREATMENT GUIDELINES

The U.S. Department of Health and Human Services (HHS) guidelines support changing HIV treatment regimens for several reasons, including:



Food requirements



Potential interactions with other drugs



Side effects

**ARE YOU READY FOR A CHANGE?**  
ASK YOUR HEALTHCARE PROVIDER IF  
*DELSTRIGO* IS RIGHT FOR YOU.

## TABLE OF CONTENTS

TAKING <i>DELSTRIGO</i>	3
OTHER MEDICATIONS	4
YOUR HEALTH AND HIV-1	5
STUDY RESULTS	5
SIDE EFFECTS	6
ABOUT CHOLESTEROL	7
QUESTIONS FOR YOUR HEALTHCARE PROVIDER	8

## IMPORTANT SAFETY INFORMATION (CONT'D)

Do not take DELSTRIGO if you are currently taking any of the following medicines:

- carbamazepine
- oxcarbazepine
- phenobarbital
- phenytoin
- enzalutamide
- rifampin
- rifapentine
- mitotane
- St. John's wort

Ask your healthcare provider or pharmacist if you are not sure if your medicine is one that is listed above. If you have taken any of the medicines in the past 4 weeks, talk to your healthcare provider or pharmacist before starting DELSTRIGO.



# TAKING DELSTRIGO

Some life changes can be an opportunity to form new habits. You and your healthcare provider can decide if DELSTRIGO could be right for you.



DELSTRIGO is **1 pill** for HIV-1 usually taken by itself (without other HIV medicines)



Take DELSTRIGO **1 time each day**, about the same time every day



DELSTRIGO can be taken **with or without food**

## IMPORTANT SAFETY INFORMATION (CONT'D)

**Do not take DELSTRIGO if you have ever had an allergic reaction to lamivudine.**

**Severe skin reactions** have happened in people treated with DELSTRIGO. Call your healthcare provider right away if you develop a rash during treatment with DELSTRIGO.

**Stop taking DELSTRIGO and get medical help right away if you develop a painful rash with any of the following symptoms:** fever, blisters or sores in the mouth, blisters or peeling of the skin, or redness or swelling of the eyes (conjunctivitis).

New or worse kidney problems, including kidney failure, can happen while you are taking DELSTRIGO. Your healthcare provider should do blood and urine tests to check your kidneys.



Models living with HIV

If you develop new or worse kidney problems, your healthcare provider may tell you to stop taking DELSTRIGO.

Some people who take DELSTRIGO experience bone problems such as pain, softening, or thinning of the bone. Tell your healthcare provider if you have any of the following symptoms during treatment with DELSTRIGO: bone pain that does not go away or worsening bone pain; pain in your arms, legs, hands or feet; broken bones; or muscle pain or weakness.

**Delstrigo**  
doravirine/lamivudine/  
tenofovir disoproxil fumarate  
100 mg/300 mg/300 mg tablets

## DELSTRIGO AND OTHER MEDICATIONS

Getting to and staying undetectable is extremely important, and it's also important to consider whether your HIV-1 treatment can be used with other medicines.

**DELSTRIGO can be taken with a wide range of non-HIV-1 medications, such as:**

Statins for **high cholesterol** (atorvastatin)

---

Metformin for **diabetes**

---

Proton pump inhibitors for **heartburn** (pantoprazole)

---

Oral contraceptives for **birth control** (ethinyl estradiol and levonorgestrel)

---

Methadone for **pain management**

*DELSTRIGO should not be taken with medicines that are toxic to the kidney (known as 'nephrotoxic') such as high dose or multiple nonsteroidal anti-inflammatory drugs (NSAIDs). DELSTRIGO should not be taken with these medicines or when these medicines have been taken recently.*

## IMPORTANT SAFETY INFORMATION (CONT'D)

Changes in your immune system (Immune Reconstitution Syndrome) can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having new symptoms after starting your HIV-1 medicine.



## IMPORTANT SAFETY INFORMATION (CONT'D)

The most common side effects of DELSTRIGO include: dizziness, nausea, and abnormal dreams.

These are not all the possible side effects of DELSTRIGO. For more information, ask your healthcare provider or pharmacist.

Before starting DELSTRIGO, tell your healthcare provider about all your medical conditions, including if you have hepatitis B virus infection; kidney problems; bone problems, including a history of bone fractures; are pregnant or plan to become pregnant. It is not known if DELSTRIGO can harm your unborn baby. Tell your healthcare provider if you are breastfeeding or plan to breastfeed. The medicines in DELSTRIGO may pass to your baby in your breast milk. HIV may pass to your baby through your breast milk if your baby does not have HIV.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. **Some medicines interact with DELSTRIGO.** Keep a list of your medicines to show your healthcare provider and pharmacist.

**Delstrigo**<sup>®</sup>  
doravirine/lamivudine/  
tenofovir disoproxil fumarate  
100 mg/300 mg/300 mg tablets

# YOUR HEALTH AND HIV-1

## STAYING UNDETECTABLE IS IMPORTANT

DELSTRIGO can help you get your viral load to undetectable. Undetectable means the level of HIV in your blood is so low that lab tests can't detect it. And when your HIV is under control, you can focus on your health and life goals beyond HIV.

In a 48-week study, adults living with HIV-1 who were undetectable on their current treatment regimen switched to **DELSTRIGO** either at the start of the study, or at week 24 of the study.

## IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider if you have taken rifabutin in the past 4 weeks. **Do not start taking a new medicine without telling your healthcare provider.** Your healthcare provider can tell you if it is safe to take DELSTRIGO with those other medicines.




### AT THE START OF THE STUDY:

- **447** adults switched to DELSTRIGO
- **91%** of those adults stayed undetectable at week 48

### AT WEEK 24 OF THE STUDY:

- **209** adults switched to DELSTRIGO
- **95%** of those adults were undetectable at week 24 before switching to DELSTRIGO
- **95%** stayed undetectable at week 48

### AT WEEK 48 OF THE STUDY:



DELSTRIGO may not have these effects on all adults living with HIV.

## MORE ABOUT THIS STUDY

Participants were undetectable (HIV-1 RNA <50 copies/mL) for at least 6 months before starting the trial and had no history of resistance\* to DELSTRIGO.

Before switching to DELSTRIGO, participants were taking one of the following treatment regimen options:

2 NRTIs in combination with a PI plus either

- ritonavir or cobicistat
- elvitegravir plus cobicistat, or
- an NNRTI

Participants were assigned by chance to one of the two treatment groups in the study. Participants and their healthcare providers taking part in the study knew which group they were part of.

**Acronyms:** NNRTI, non-nucleoside reverse transcriptase inhibitor; NRTI, nucleoside reverse transcriptase inhibitor; PI, protease inhibitor

\*Resistance happens when the virus changes form and your medicine stops working.

**Delstrigo**  
doravirine/lamivudine/  
tenofovir disoproxil fumarate  
100 mg/300 mg/300 mg tablets






## A LOW RATE OF SIDE EFFECTS

In a 96-week study of adults who had never been on an HIV-1 treatment before, 364 adults received **DELSTRIGO** while 364 adults received an *Atripla* (EFV/FTC/TDF) once-daily tablet.

Fewer adults stopped taking DELSTRIGO than *Atripla* because of side effects.



In this study, the most common side effects of DELSTRIGO were dizziness **7%**, nausea **5%**, and abnormal dreams **5%**.

**LIFE CHANGES.  
STAY INSPIRED.**

### IMPORTANT SAFETY INFORMATION (CONT'D)

Take DELSTRIGO every day exactly as your healthcare provider tells you to take it. Take DELSTRIGO 1 time each day, at about the same time every day.

DELSTRIGO is usually taken by itself (without other HIV medicines).

If you take the medicine rifabutin during treatment with DELSTRIGO, your healthcare provider will also prescribe an additional dose of doravirine for you.

### IMPORTANT SAFETY INFORMATION (CONT'D)

Carefully follow your healthcare provider's instruction about when to take doravirine and how much to take. This is usually 1 tablet of doravirine about 12 hours after your last dose of DELSTRIGO. Do not change your dose or stop taking DELSTRIGO without talking to your healthcare provider first.


## ABOUT CHOLESTEROL

HIV-1 and/or treatments for HIV may have a negative effect on your “bad” cholesterol (LDL-C).

In a 96-week study of adults who were new to treatment, 320 adults received DELSTRIGO and 307 adults received *Atripla*.

DELSTRIGO did not increase “bad” cholesterol, while *Atripla* did at week 48.

**At week 48, the effects on “bad” cholesterol (LDL-C) were:**



The effects on “bad” cholesterol at week 96 were similar to those seen at week 48.

**The benefit of these changes has not been demonstrated.**

### IMPORTANT SAFETY INFORMATION (CONT'D)

Take DELSTRIGO on a regular dosing schedule as instructed by your healthcare provider. Do not miss doses.

Get your DELSTRIGO refilled from your healthcare provider or pharmacy before you run out.



*When life brings change,  
what centers me?*

**MY HOME  
BASE**

**Delstrigo**   
doravirine/lamivudine/  
tenofovir disoproxil fumarate  
100 mg/300 mg/300 mg tablets



# STARTING THE CONVERSATION

Only you and your healthcare provider know what treatment is right for you. Here are some questions you can use to start a conversation about your treatment options.



How will my treatment fit into my routine?



What side effects could I experience?



Could my HIV-1 treatment impact my weight?



Can my HIV-1 treatment have a negative effect on my cholesterol?



Can I still take other medications with my HIV-1 medication?



Do I need to eat when I take my HIV-1 medication?



Are there other health risks my HIV or HIV-1 treatment could cause?



Model living with HIV



## AN ADDITIONAL TREATMENT CONSIDERATION

Maintaining a healthy weight is an important part of taking care of your health and managing your HIV. You may have concerns about weight change. If you do, talk to your healthcare provider.

## IMPORTANT SAFETY INFORMATION

### Worsening of hepatitis B virus infection (HBV).

If you have both Human Immunodeficiency Virus-1 (HIV-1) and HBV and stop taking DELSTRIGO, your HBV may suddenly get worse. Do not stop taking DELSTRIGO without first talking to your healthcare provider, as they will need to monitor your health. Your healthcare provider should test you for HBV infection before you start treatment with DELSTRIGO.

Please read the accompanying [Patient Information](#) for DELSTRIGO, including an important warning about the potential worsening of hepatitis B after stopping DELSTRIGO, and discuss it with your healthcare provider. The physician [Prescribing Information](#) also is available.



TAKE A PIC OF THESE QUESTIONS AND DISCUSS THEM WITH YOUR HEALTHCARE PROVIDER.





*When life brings change,  
what keeps me connected?*

# COMMUNITY

**LIFE CHANGES.  
STAY CONNECTED.**

**ASK YOUR HEALTHCARE  
PROVIDER IF DELSTRIGO IS  
RIGHT FOR YOU.**

**READY TO LEARN MORE  
ABOUT DELSTRIGO?**



Scan this code and find out if  
DELSTRIGO is right for you,  
or go to [delstrigo.com](http://delstrigo.com).





## FREE TRIAL OFFER FOR *DELSTRIGO*

Eligible patients may receive a free 30-tablet trial supply of *DELSTRIGO*



Scan the QR code to learn about  
a Free Trial Offer for *DELSTRIGO*.

Not all patients are eligible. Certain restrictions apply. Please see Terms and Conditions.



Visit [delstrigo.com/special-offers](https://delstrigo.com/special-offers) to determine if you are eligible for the Free Trial Offer for *DELSTRIGO*.



Save the voucher to your mobile device or print it.



Present the voucher with a valid signed prescription to any participating eligible retail pharmacy (certain restrictions apply).

There is no requirement to purchase any product or service for you to receive a free 30-tablet trial supply of *DELSTRIGO*. Not all patients are eligible. Certain restrictions apply. See Terms and Conditions.

Please read the accompanying [Patient Information](#) for *DELSTRIGO*, including an important warning about the potential worsening of hepatitis B after stopping *DELSTRIGO*, and discuss it with your healthcare provider. The physician [Prescribing Information](#) also is available.

Products mentioned are trademarks of their respective owners.



Copyright © 2025 Merck & Co., Inc., Rahway, NJ, USA and its affiliates.  
All rights reserved. US-DOR-01467 03/25

**Delstrigo**<sup>®</sup>  
doravirine/lamivudine/  
tenofovir disoproxil fumarate  
100 mg/300 mg/300 mg tablets